A IOURNAL OF NEUROLOGY

Neural correlates of bimanual anti-phase and in-phase movements in Parkinson's disease

Tao Wu,¹ Liang Wang,² Mark Hallett,³ Kuncheng Li² and Piu Chan¹

- 2 Department of Radiology, Xuanwu Hospital, Capital Medical University, Beijing, 20892-1428, People's Republic of China
- 3 Human Motor Control Section, Medical Neurology Branch, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892-1428, USA

Correspondence to: Tao Wu, Department of Neurobiology, Key Laboratory on Neurodegenerative Disorders of Ministry of Education, Beijing Institute of Geriatrics, Xuanwu Hospital, Capital Medical University, 45 Changchun Street, Beijing, 100053, People's Republic of China E-mail: wutao69@gmail.com

Patients with Parkinson's disease have great difficulty in performing bimanual movements; this problem is more obvious when they perform bimanual anti-phase movements. The underlying mechanism of this problem remains unclear. In the current study, we used functional magnetic resonance imaging to study the bimanual coordination associated changes of brain activity and inter-regional interactions in Parkinson's disease. Subjects were asked to perform right-handed, bimanual in-phase and bimanual anti-phase movements. After practice, normal subjects performed all tasks correctly. Patients with Parkinson's disease performed in-phase movements correctly. However, some patients still made infrequent errors during anti-phase movements; they tended to revert to in-phase movement. Functional magnetic resonance imaging results showed that the supplementary motor area was more activated during anti-phase movement than in-phase movement in controls, but not in patients. In performing anti-phase movements, patients with Parkinson's disease showed less activity in the basal ganglia and supplementary motor area, and had more activation in the primary motor cortex, premotor cortex, inferior frontal gyrus, precuneus and cerebellum compared with normal subjects. The basal ganglia and dorsolateral prefrontal cortex were less connected with the supplementary motor area, whereas the primary motor cortex, parietal cortex, precuneus and cerebellum were more strongly connected with the supplementary motor area in patients with Parkinson's disease than in controls. Our findings suggest that dysfunction of the supplementary motor area and basal ganglia, abnormal interactions of brain networks and disrupted attentional networks are probably important reasons contributing to the difficulty of the patients in performing bimanual anti-phase movements. The patients require more brain activity and stronger connectivity in some brain regions to compensate for dysfunction of the supplementary motor area and basal ganglia in order to perform bimanual movements correctly.

Keywords: Parkinson's disease; bimanual movements; fMRI; brain activity; effective connectivity

Abbreviations: fMRI = functional magnetic resonance imaging; FWE = family-wise error; PPI = psychophysiological interaction; SM1 = primary sensorimotor cortex; SMA = supplementary motor area; UPDRS = Unified Parkinson's Disease Rating Scale

¹ Key Laboratory on Neurodegenerative Disorders of Ministry of Education, Department of Neurobiology, Beijing Institute of Geriatrics, Xuanwu Hospital, Capital Medical University, Beijing 100053, People's Republic of China

Received January 5, 2010. Revised April 29, 2010. Accepted April 30, 2010. Advance Access publication June 20, 2010 © The Author (2010). Published by Oxford University Press on behalf of the Guarantors of Brain. All rights reserved. For Permissions, please email: journals.permissions@oxfordjournals.org

Introduction

Bimanual movements are important to daily life and require temporal and spatial coordination. Patients with Parkinson's disease commonly show impaired bimanual coordination. This problem is more obvious when they perform bimanual anti-phase movements than in-phase movements (Johnson et al., 1998; Serrien et al., 2000; van den Berg et al., 2000; Geuze, 2001; Almeida et al., 2002; Ponsen et al., 2006). Anti-phase bimanual movements, for example at the wrist, occur when both hands perform the same movement, but with a phase shift of 180° between the two hands (i.e. one hand flexes while the other extends). For in-phase bimanual wrist movements, the two hands move in and out together, requiring simultaneous flexion and extension at both wrists, without phase shift. Both in-phase and anti-phase movements require synchronization between the two hands, but the anti-phase movements additionally need contralateral movement suppression (of a mirrored movement) and the independence of the two movements. Transcranial magnetic stimulation during a bimanual in-phase task could simultaneously reset the rhythmic movements of both hands. In contrast, the transcranial magnetic stimulation has little effect on the bimanual anti-phase task, indicating that control of rhythm differs in the anti- and in-phase tasks (Chen et al., 2005). In-phase rhythms are more stable than anti-phase rhythms (Tuller and Kelso, 1989) and are easier to perform than anti-phase movements.

It has been shown that patients with Parkinson's disease can perform bimanual in-phase movements correctly, but perform anti-phase movements with more error and variability (Johnson *et al.*, 1998; Almeida *et al.*, 2002). Patients have a tendency to revert anti- to in-phase movements (Johnson *et al.*, 1998). Additionally, the performance of anti-phase movements is not improved with the presence of external pacing cues (Johnson *et al.*, 1998; Almeida *et al.*, 2002). The difficulty in performing bimanual movements, especially in performing anti-phase movements, can be detected even in early Parkinson's disease (Ponsen *et al.*, 2006). Therefore, understanding the neural mechanism of this problem is not only useful for our understanding about the pathophysiology of movement in Parkinson's disease, but also may help to develop a sensitive clinical procedure to assess and quantify the illness (Johnson *et al.*, 1998; Ponsen *et al.*, 2006).

The aim of the current study was to use functional magnetic resonance imaging (fMRI) to explore brain activations, as well as interactions within brain networks during performance of bimanual movements in patients with Parkinson's disease. Previous studies have shown that the control of bimanual coordination cannot be assigned to a single area; rather, it seems to involve a distributed network in which interactive processes take place between many neural assemblies to ensure efferent organization and sensory integration (Debaere *et al.*, 2001; Swinnen, 2002). Thus, investigations about interactions among brain regions may play a more important role than simply exploring activity in understanding bimanual movement-related brain functional changes. The methods used to explore inter-regional interactions in a given task are analysis of functional connectivity (Friston *et al.*, 1993*b*). These methods are

increasingly being used to investigate Parkinson's disease induced modifications of brain networks (Rowe *et al.*, 2002; Ma and Wang, 2008; Helmich *et al.*, 2009; Palmer *et al.*, 2009; van Eimeren *et al.*, 2009; Wu *et al.*, 2010). In healthy subjects, the supplementary motor area (SMA) has been suggested to be critical for bimanual coordination (Sadato *et al.*, 1997*b*; Stephan *et al.*, 1999; Toyokura *et al.*, 1999; Immisch *et al.*, 2001). In addition, impaired activity in the SMA is a common finding in Parkinson's disease (Playford *et al.*, 1992; Jahanshahi *et al.*, 1995; Rascol *et al.*, 1997; Haslinger *et al.*, 2001; Buhmann *et al.*, 2003). Thus, we investigated the effective connectivity in the SMA to explore Parkinson's disease-related changes in interactions of neural networks in bimanual movements.

Methods

Subjects

We studied 15 patients with Parkinson's disease, aged 44-71 years (mean 59.73 years) and included 10 males and 5 females. The diagnosis of Parkinson's disease was based on medical history, physical and neurological examinations, response to L-dopa and laboratory tests and MRI scans to exclude other diseases. Patients were assessed with the Unified Parkinson's Disease Rating Scale (UPDRS) (Lang and Fahn, 1989), the Hoehn and Yahr disability scale (Hoehn and Yahr, 1967) and Mini-Mental State Exam while off their medications. The Mini-Mental State Exam was \geq 27 in all subjects and there was no difference between the patients and controls. The clinical data are shown in Table 1. We also investigated 15 age- and sex-matched normal subjects (aged 44-73 years, mean 60.30) as controls. All subjects were right handed as measured by the Edinburgh Inventory (Oldfield, 1971). The experiments were performed according to the Declaration of Helsinki and were approved by the Institutional Review Board. All subjects gave their written informed consent for the study.

Experimental design

Subjects were asked to perform three types of finger movements: (i) extension and flexion of the right index finger (unimanual right-hand movement); (ii) extension and flexion of both index fingers simultaneously (bimanual in-phase movement); and (iii) simultaneous extension of one index finger and flexion of the other index finger, and vice versa, to produce movement in the same spatial direction (bimanual anti-phase movement). All movements were self-paced and were executed at an interval of 2 s. No external cue was given to help the subjects move at the specified rate. Movement amplitude was determined as the maximal possible for both extension and

Table 1	Clinical	details	of	patients	with	Parkinson's
disease	(mean \pm	SD)				

Age (years)	59.73 ± 8.27
Sex	5 female, 10 male
Disease duration (years)	3.47 ± 1.60
UPDRS motor score (off medication)	20.67 ± 3.48
Hoehn and Yahr staging (off medication)	1.70 ± 0.37
∟-dopa dose (mg/day)	333.33 ± 48.80

flexion. In the flexion direction, it was limited by a response device fixed to their hand. Before the fMRI, all subjects practiced until they could perform all the tasks properly and move at the required rate.

Functional MRI procedure

Patients were scanned only after their medication had been withdrawn for at least 12 h. Imaging was performed on a 1.5 T Siemens Sonata scanner. High-resolution axial T_1 - and T_2 -weighted images were obtained in every subject to exclude other neurological disorders. We used an echo planar imaging gradient sequence sensitive enough to acquire functional images (repetition time = 2000 ms. echo time = 60 ms, flip angle = 90° , field of view = 24×24 cm, matrix = 64×64). Twenty axial slices were collected with 5 mm thickness and a 2 mm gap. We had three fMRI sessions and the three movement tasks were performed randomly once during each scan. Two conditions were contained in each scanning session and were defined as the 'rest' and 'active' conditions, respectively. Each condition lasted 20s and was repeated six times in a session. During the rest condition, subjects were instructed to keep their eyes closed and to remain motionless. The active condition in each session contained one motor task. Each subject's performance during fMRI for each task was monitored by an investigator and recorded by video. If there were any errors of finger movement, we asked the subject to repeat that session until he/she could perform it correctly. Additionally, two response devices were fixed to each of their hands to record the rate of movements during fMRI scanning.

Data analysis

Behavioural data analysis

Each subject's performance for each task was recorded and compared between the patients and normal subjects (two-sample *t*-test, P < 0.05). Additionally, the frequencies of the movements were compared between the groups.

Brain activity analysis

Functional MRI data were analysed with Statistical Parametric Mapping 2 software (Wellcome Institute of Cognitive Neurology, London, UK). They were slice-time corrected and aligned to the first image of each session for motion correction. After spatial normalization, all images were resampled into voxels that were $2 \times 2 \times 2$ mm in size and smoothed with a Gaussian filter of 6mm full-width at half maximum. In the first-level, data were analysed for each subject separately on a voxel-by-voxel basis using the general linear model approach for the time series. We defined a model using a fixed effect boxcar design convolved with a haemodynamic response function to analysis of task-dependent activation. A contrast representing the effect of the active condition compared with the rest condition was defined and contrast images were calculated individually for each condition. These contrast images were used in the second-level for random effects analyses. For the within group analysis, a one-sample t-test model was used to identify the brain activity for each task [P < 0.05, family-wise error (FWE) corrected]. Then, a paired *t*-test was used to compare the results between anti-phase and right-hand movement, in-phase and right-hand movement, as well as between anti-phase and in-phase movement (P<0.05, FWE corrected). For the between-group comparisons, a two-sample t-test (P < 0.05, FWE corrected) was used to explore the difference between patients and normal subjects in performing bimanual movements.

Finally, in order to explore whether the changes of brain activity relate to disease severity, a correlation analysis of activations during bimanual movements versus the UPDRS motor score was performed in patients.

Effective connectivity analysis

Effective connectivity was assessed using the method of psychophysiological interaction (PPI) (Friston *et al.*, 1997). PPI is defined as the change in contribution of one brain area to another due to a change in experimental condition or psychological context (Friston *et al.*, 1997). It aims to explain regionally specific responses in terms of the interaction between the psychological variable and the activity in a specific index area. The analysis was constructed to test for differences in the regression slope of the activity in all remaining brain areas on the activity in the index area depending on the movement type.

Given the critical role of the SMA in bimanual coordination and defective function of this region in Parkinson's disease, we chose this region as the index area for PPI analysis. The SMA contains two separate areas: the SMA-proper in the caudal portion and the pre-SMA in the rostral portion (Tanji and Hoshi, 2001). From previous reports, the region showing stronger activation during bimanual anti-phase than bimanual in-phase movements is the SMA-proper (Sadato et al., 1997b; Toyokura et al., 1999; 2002; Immisch et al., 2001). A study compared the activation of pre-SMA and SMA-proper during unimanual and bimanual movements and found that the SMA-proper was more activated during bimanual movements than unimanual movements, whereas the pre-SMA was inconsistently activated (Toyokura et al., 2002). Thus, the index volume in the current study was defined as centred on the voxel that showed the maximum magnitude of activation within the SMA-proper with a radius of 5 mm individually for each bimanual movement. The PPI term (referred to as 'PPI regressor') was computed as the element-by-element product of the deconvolved extracted time series of the SMA-proper and a vector coding for the main effect of task (Gitelman et al., 2003; Stephan et al., 2003; Garraux et al., 2005; Wu et al., 2010). For each subject, the PPI regressor, the task regressor (representing mode of bimanual movements) and the extracted time series were entered in a first-level model of effective connectivity in which the PPI regressor was orthogonalized with regard to the main effect of the task and the regional time series. Brain areas receiving context-dependent influences from the SMA-proper were determined by testing for positive slopes of the PPI regressor. Contrast images from the first-level PPI analysis for each bimanual task in each subject were entered into a second-level, random-effect model. A one-sample t-test was used to identify the connectivity for each bimanual task for each group (P < 0.05, FWE corrected). A paired *t*-test was used to compare the results between anti- and in-phase movement in each group (P < 0.05, FWE corrected). Then, a two-sample *t*-test model was used to explore the difference between patients and controls in performing each bimanual task (P<0.05, FWE corrected).

Results

Task performance

All subjects had no obvious difficulty in performing unimanual and in-phase movements and only needed brief practice to perform the tasks without error. Both groups had more difficulty during the practice of anti-phase movements. Although the patients spent significantly more time than normal subjects in practice (mean 21.3 ± 5.2 min versus 10.6 ± 3.8 min), in the end all of them could perform the anti-phase task correctly. During fMRI scanning, the normal subjects performed all tasks without any error. All patients performed unimanual and in-phase movements correctly. In contrast, while performing anti-phase movement, although there was no formal significant difference of performance between the groups, four patients inverted a few moves to in-phase movements ($1.7\pm3.3\%$ errors, all in anti-phase movements), but recognized and corrected the errors by themselves. We asked these four patients to perform the anti-phase movement again and all of them performed the task without any error during the second session. We used the repeat performance for fMRI data analysis.

There was no between- or within-group difference for the rate of performance of motor tasks [repeated measures analysis of variance (ANOVA), P > 0.05]. In patients, the rates of movements were 0.56 ± 0.09 Hz for right-hand movement, 0.52 ± 0.11 Hz for in-phase movement and 0.52 ± 0.08 Hz for anti-phase movement. In normal subjects, the rates were 0.55 ± 0.06 Hz, 0.54 ± 0.03 Hz and 0.52 ± 0.06 Hz, respectively.

Brain activity

Within-group analysis

During performance of right-hand movements, normal subjects activated the left primary sensorimotor cortex (SM1), right premotor cortex, SMA-proper, right dorsolateral prefrontal cortex, right inferior frontal gyrus, bilateral insula, bilateral basal ganglia and bilateral cerebellum (Fig. 1A, left column; one-sample *t*-test, P < 0.05, FWE corrected). With a more liberal threshold (P < 0.05, false discovery rate corrected), the left premotor cortex was also activated. While performing both bimanual in- and anti-phase movements, the bilateral SM1, bilateral premotor cortex, right dorsolateral prefrontal cortex, left inferior frontal gyrus, SMA-proper, cingulate motor area, bilateral inferior and superior parietal lobule, precuneus, bilateral superior temporal gyrus, bilateral thalamus, bilateral basal ganglia and bilateral cerebellum were activated (Fig. 1B and C, left column; one-sample *t*-test, P < 0.05, FWE corrected).

During performance of right-hand movements, patients with Parkinson's disease activated the left SM1, left premotor cortex,



Figure 1 Brain regions activated during performing motor tasks in normal control group (left column), and in patients with Parkinson's disease group (right column). Results were thresholded at P < 0.05 (FWE corrected). (**A**) Brain areas activated during performing right-hand movements. (**B**) Brain areas activated during performing bimanual in-phase movements. (**C**) Brain areas activated during performing bimanual anti-phase movements.

SMA-proper, left inferior frontal gyrus, bilateral insula, left putamen and bilateral cerebellum (Fig. 1A, right column; one-sample *t*-test, P < 0.05, corrected). While performing both bimanual in- and anti-phase movements, the bilateral SM1, bilateral premotor cortex, right dorsolateral prefrontal cortex, SMA-proper, cingulate motor area, bilateral inferior and superior parietal lobule, precuneus, bilateral superior temporal gyrus, left thalamus, bilateral basal ganglia and bilateral cerebellum were activated (Fig. 1B and C, left column; one-sample *t*-test, P < 0.05, FWE corrected).

In patients with Parkinson's disease performing bimanual antior in-phase movements, there was more activation in the right SM1, left inferior frontal gyrus, right premotor cortex, bilateral superior parietal lobule, SMA-proper, cingulate motor area, precuneus and left cerebellum compared to right-hand movement alone. In normal subjects, in addition to these regions, the right dorsolateral prefrontal cortex, bilateral thalamus, right putamen and right globus pallidus were more activated in performing bimanual anti- or in-phase movements compared to right-hand movements (Fig. 2A; paired t-test, P<0.05, FWE corrected). In controls, the SMA-proper, bilateral premotor cortex, left inferior frontal gyrus, right post-central gyrus, left inferior parietal lobule and bilateral cerebellum were more activated for anti-phase movements than for in-phase movements. In patients, performing anti-phase movements utilized more activation in the bilateral inferior frontal gyrus, left middle frontal gyrus, bilateral premotor cortex, right precentral gyrus, bilateral post-central gyrus, left inferior temporal gyrus and bilateral cerebellum than in performing in-phase movements (Table 2 and Fig. 2B; paired t-test, P < 0.05, FWE corrected).

Between-group comparisons

During the performance of in-phase movements, patients with Parkinson's diseasehad greater activity in the right SM1, left premotor cortex, bilateral post-central gyrus, left superior parietal lobule, right precuneus and bilateral cerebellum, and had less activity in the SMA-proper, bilateral thalamus, left putamen and right globus pallidus compared with normal subjects (Table 3; two sample *t*-test, P < 0.05, FWE corrected).

During the performance of anti-phase movements, patients had more activation in the left SM1, left premotor cortex, right inferior frontal gyrus, bilateral precentral gyrus, bilateral postcentral gyrus, left superior parietal lobule, left inferior parietal lobule, bilateral paracentral lobule, bilateral precuneus and bilateral cerebellum, and less activity in the SMA-proper, bilateral thalamus and right globus pallidus compared with normal subjects (Fig. 3 and Table 4; two sample *t*-test, P<0.05, FWE corrected). With a more liberal threshold (P<0.001, uncorrected), we also found that the left SM1 was more activated during in-phase movements, whereas the right SM1 was more activated during anti-phase movements in patients compared with normal controls.

Correlation analysis

A correlation analysis found that during in-phase movement, brain activations in the left putamen and SMA-proper were negatively correlated with the UPDRS motor score, whereas the activations in the right inferior frontal gyrus, right inferior parietal lobule, bilateral precuneus and bilateral cerebellum were positively correlated with the UPDRS motor score (P<0.05, FWE corrected).



Figure 2 (A) Brain areas more activated for bimanual anti-phase movements than for right-hand movements in normal control group (left column) and in patients with Parkinson's disease group (right column). (B) Brain areas more activated for bimanual anti-phase movements than for bimanual in-phase movements in normal control group (left column) and in patients with Parkinson's disease (right column). Results were thresholded at P < 0.05 (FWE corrected).

Table 2 Brain areas more activated in performing anti-phase movements than in performing in-phase movements in normal subjects and patients with Parkinson's disease

Brain region	Coordinates			t-value	Cluster size
	x	у	z		
Normal subjects					
Right cerebellum, anterior lobe, culmen	10	-46	-16	10.28	698
Left inferior parietal lobule	-50	-30	31	9.62	231
Left inferior frontal gyrus	-58	13	23	9.35	194
Left premotor cortex	-16	-5	54	9.33	119
SMA-proper	0	-2	64	9.32	317
Left cerebellum, posterior lobe, uvula	-6	-81	-33	9.05	202
Right post-central gyrus	36	-36	64	8.71	347
Right premotor cortex	24	-6	42	8.69	303
Parkinson's disease patients					
Right inferior frontal gyrus	20	27	-3	10.09	161
Left middle frontal gyrus	-34	36	17	9.40	203
Left inferior frontal gyrus	-36	13	-14	9.11	103
Left cerebellum, posterior lobe, tuber	-36	-66	-30	9.06	200
Right post-central gyrus	26	-49	67	8.99	361
Left post-central gyrus	-24	-37	68	8.59	102
Left premotor cortex	-12	-2	65	8.26	206
Right cerebellum, posterior lobe, declive	48	-51	-18	8.17	196
Right premotor cortex	24	-2	58	8.15	216
Right precentral gyrus	51	12	5	8.10	164

List of the brain regions showing significantly more activity in performing anti-phase movements than in performing in-phase movements in each group (paired *t*-test, P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).

Table 3 Differences of brain activity between patients with Parkinson's disease and normal subjects in performing in-phase movements

Brain region	Coordinates			t-value	Cluster size
	x	у	z		
Normal–Parkinson's disease					
Left thalamus	-2	-8	-3	9.17	182
SMA-proper	2	-8	62	9.09	84
Left putamen	-20	-2	17	8.71	59
Right thalamus	18	-16	-2	8.37	87
Right globus pallidus	16	-8	-4	8.09	57
Parkinson's disease–normal					
Left post-central gyrus	-42	-22	32	10.29	511
Left superior parietal lobule	-28	-50	56	9.07	221
Right cerebellum, posterior lobe, tonsil	30	-42	-32	8.69	40
Right precuneus	24	-48	48	8.64	65
Right post-central gyrus	53	-21	53	8.41	24
Right SM1	14	-26	66	8.25	54
Right cerebellum, posterior lobe, tonsil	16	-54	-34	8.21	31
Left premotor cortex	-18	-9	54	8.21	34
Right cerebellum, anterior lobe, culmen	2	-36	-23	8.10	22
Left cerebellum, posterior lobe, tonsil	-36	-50	-38	8.06	33

List of the brain regions showing significantly more activity in normal subjects than in patients with Parkinson's disease (Normal–Parkinson's disease), or more activity in patients with Parkinson's disease than in normal subjects (Parkinson's disease–Normal), in performing in-phase movements (two sample *t*-test, P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).

A correlation analysis on anti-phase movement found that brain activations in the bilateral putamen and SMA-proper were negatively correlated with the UPDRS motor score, whereas the activity in the right inferior frontal gyrus, bilateral premotor cortex, bilateral inferior parietal lobule, bilateral precuneus and bilateral cerebellum was positively correlated the UPDRS motor score (P < 0.05, FWE corrected). In this study, negative correlation means that as the UPDRS motor score increased, the brain



Figure 3 Brain areas more activated in normal subjects than in patients with Parkinson's disease (**A**) and more activated in patients with Parkinson's disease than in normal subjects during performing anti-phase movements (**B**). Results were thresholded at P < 0.05 (FWE corrected).

activations are weaker, and positive correlation means that as the UPDRS motor score increased, the brain activations are stronger.

Effective connectivity analysis

Within-group analysis

PPI analysis found that in normal subjects during in-phase movement, the SMA-proper had significant connections with the bilateral SM1, right premotor cortex, right insula, right middle frontal gyrus, right globus pallidus, right putamen, right subthalamic nucleus, right substania nigra, left parahippocampal gyrus and bilateral cerebellum (Fig. 4A, left column; one-sample *t*-test, P < 0.05, FWE corrected). In performing anti-phase movements, normal subjects had the bilateral SM1, SMA-proper, pre-SMA, bilateral premotor cortex, left dorsolateral prefrontal cortex, left cingulate motor area, left temporal lobe, left precuneus, left insula, right thalamus, left putamen, right globus pallidus and bilateral cerebellum effectively connected with the SMA-proper (Fig. 4A, right column; one-sample *t*-test, P < 0.05, FWE corrected).

In patients with Parkinson's disease, the SMA-proper was connected with the bilateral SM1, right post-central gyrus, left inferior parietal lobule, right angular gyrus, left precuneus, right globus pallidus, right thalamus and bilateral cerebellum during performance of in-phase movements (Fig. 4B, left column; one-sample *t*-test, *P* < 0.05, FWE corrected). In performing anti-phase movements, patients had the bilateral SM1, right postcentral gyrus, SMA-proper, right premotor cortex, right insula, bilateral superior parietal lobule, bilateral inferior parietal lobule, left inferior temporal gyrus, right globus pallidus, right putamen, right thalamus and bilateral cerebellum connected with the SMA-proper (Fig. 4B, right column; one-sample *t*-test, *P* < 0.05, FWE corrected).

In normal subjects, the SMA-proper was more connected with the left SM1, right premotor cortex, left dorsolateral prefrontal cortex, left cingulate motor area, left precuneus, right limbic lobe, left putamen and bilateral cerebellum during anti-phase movement than in-phase movement (Table 5 and Fig. 5A; paired *t*-test, P < 0.05, FWE corrected). The SMA-proper was more connected with the right SM1 (x = 30, y = -20, z = 64, cluster size 56) in performing in-phase movements than in performing anti-phase movements (paired *t*-test, P < 0.05, FWE corrected).

In patients, the SMA-proper was more connected with the left SM1, right precentral gyrus, right premotor cortex, right post-central gyrus, bilateral inferior parietal lobule, left paracentral lobule and right precuneus during anti-phase movement than in-phase movement (Table 5 and Fig. 5B; paired *t*-test, P < 0.05, FWE corrected). The SMA-proper was more connected with the right SM1 (x = 32, y = -20, z = 70, cluster size 87) in performing in-phase movements than in performing anti-phase movements (paired *t*-test, P < 0.05, FWE corrected).

Between-group comparisons

During the performance of in-phase movements, patients with Parkinson's disease had more connectivity to the left SM1, left post-central gyrus, left precuneus and bilateral cerebellum with the SMA-proper compared with normal controls (Table 6 and Fig. 6A, upper row; paired *t*-test, P < 0.05, FWE corrected). Normal subjects showed more connectivity in the right premotor cortex and right putamen compared to patients (Table 6, Fig. 6A, lower row; paired *t*-test, P < 0.05, FWE corrected).

During the performance of anti-phase movements, patients had more connectivity to the bilateral SM1, bilateral superior parietal lobule, right precuneus and bilateral cerebellum with the SMA-proper compared with controls (Table 7 and Fig. 6B, upper row; paired *t*-test, P < 0.05, FWE corrected). Normal subjects showed more connectivity to the left dorsolateral prefrontal cortex and left putamen compared to patients (Table 7 and Fig. 6B, lower row; paired *t*-test, P < 0.05, FWE corrected).

Discussion

The present study, for the first time, explored the neural mechanisms underlying the difficulty in performing bimanual tasks in Parkinson's disease. The novel findings are that the patterns of brain activity, as well as interactions of brain networks, are changed in patients with Parkinson's disease, revealing how dysfunction of the basal ganglia influences the rest of the brain.

Table 4 Differences of brain activity between patients with Parkinson's disease and normal subjects in performing anti-phase movements

Brain region	Coordinates			t-value	Cluster size
	x	у	Z		
Normal–Parkinson's disease					
SMA-proper	4	-6	54	8.86	136
Right thalamus	14	-10	2	8.59	106
Right globus pallidus	20	-14	-1	8.48	42
Left thalamus	-14	-12	0	8.16	31
Parkinson's disease-Normal					
Left precentral gyrus	-54	-3	9	9.46	323
Left cerebellum, posterior lobe, tonsil	-8	-45	-40	8.77	55
Left inferior parietal lobule	-51	-30	29	8.55	182
Left post-central gyrus	-36	-44	61	8.33	170
Left superior parietal lobule	-34	-46	50	8.14	114
Left precuneus	-16	-44	46	8.11	212
Right cerebellum, posterior lobe, declive	38	-67	-20	7.91	63
Right precentral gyrus	63	5	18	7.74	90
Right inferior frontal gyrus	46	-3	18	7.67	133
Right cerebellum, posterior lobe, tonsil	14	-56	-36	7.49	79
Left post-central gyrus	-14	-53	65	7.46	59
Left SM1	-28	-22	67	7.40	160
Left premotor cortex	-18	-14	62	7.35	84
Left paracentral lobule	-8	-39	68	7.34	120
Right paracentral lobule	6	-34	68	7.06	62
Right precuneus	2	-36	46	7.04	53

List of the brain regions showing significantly more activity in normal subjects than in patients with Parkinson's disease (Normal–Parkinson's disease), or more activity in patients with Parkinson's disease than in normal subjects (Parkinson's disease–Normal), in performing anti-phase movements (two sample *t*-test, P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).



Figure 4 Brain regions effectively connected with the SMA-proper during performing in-phase (left column) and anti-phase movements (right column), in (**A**) normal subjects and (**B**) patients with Parkinson's disease. Results were thresholded at P < 0.05 (FWE corrected) and rendered over a standard anatomical brain.

 Table 5
 Brain areas stronger connected with the SMA in the anti-phase state compared to the in-phase state in normal subjects and patients with Parkinson's disease

Brain region	Coordinates			t-value	Cluster size
	x	у	z		
Normal subjects					
Left cerebellum, anterior lobe	-2	-35	-32	10.44	86
Right cerebellum, posterior lobe, tonsil	38	-64	-32	10.12	411
Left precuneus	-14	-50	41	9.85	156
Right cerebellum, anterior lobe, culmen	14	-54	-2	9.82	63
Left cerebellum, posterior lobe, pyramis	-40	-77	-33	9.71	217
Left putamen	-24	-6	6	9.60	142
Left cingulate motor area	-16	13	32	9.41	52
Left SM1	-38	-23	42	9.36	89
Right cerebellum, posterior lobe, pyramis	18	-62	-27	9.36	42
Left dorsolateral prefrontal cortex	-26	31	35	9.10	69
Left cerebellum, anterior lobe, culmen	-40	-56	-28	9.04	36
Right limbic lobe	20	-66	11	8.91	45
Right premotor cortex	20	-9	58	8.84	21
Patients with Parkinson's disease					
Right precentral gyrus	55	-10	32	9.98	76
Left SM1	-18	-30	68	9.44	168
Left inferior parietal lobule	-42	-32	33	9.07	64
Right inferior parietal lobule	36	-42	40	8.76	51
Left paracentral lobule	-20	-42	54	8.57	38
Right precuneus	30	-44	50	8.47	26
Right premotor cortex	22	-11	46	8.42	22
Right post-central gyrus	26	-34	50	8.41	38

List of the brain regions showing a significant connectivity with the SMA (P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).



Figure 5 Brain areas more connected with the SMA-proper for bimanual anti-phase movements than for in-phase movements in normal subjects (**A**) and in patients with Parkinson's disease (**B**). Results were thresholded at P < 0.05 (FWE corrected).

The abnormal neural activity appears to explain the difficulty of bimanual coordination in Parkinson's disease.

After practice, although a few patients still made infrequent errors while performing anti-phase movements, all patients could

perform the bimanual anti- and in-phase movements correctly. The significantly more time needed for practice, and more errors made, demonstrated that patients with Parkinson's disease had more difficulty in performing bimanual movements, especially

Brain region	Coordinates			t-value	Cluster size
	x	у	z		
Normal–Parkinson's disease					
Right putamen	26	-6	4	9.46	79
Right premotor cortex	20	-6	54	8.48	25
Parkinson's disease–Normal					
Right cerebellum, posterior lobe, tonsil	24	-62	-32	9.97	81
Left cerebellum, posterior lobe, vermis	-2	-74	-36	9.83	32
Left cerebellum, posterior lobe, pyramis	-18	-68	-30	9.43	54
Left post-central gyrus	-32	-29	38	9.27	106
Left cerebellum, posterior lobe, tuber	-30	-79	-30	9.07	97
Right cerebellum, posterior lobe, declive	-30	-57	54	8.88	24
Left precuneus	-14	-50	41	8.76	82
Left SM1	-46	-17	36	8.72	31

 Table 6
 Differences of effective connectivity in the SMA between patients with Parkinson's disease and normal subjects during performing in-phase movements

List of the brain regions showing a significant connectivity with the SMA (P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).

Table 7 Differences of effective connectivity in the SMA between patients with Parkinson's disease and normal subjects during performing anti-phase movements

Brain region	Coordinates			t-value	Cluster size
	x	у	z		
Normal–Parkinson's disease					
Left dorsolateral prefrontal cortex	-30	26	34	9.56	323
Left Putamen	-26	-8	0	8.69	118
Parkinson's disease–Normal					
Left SM1	-32	-30	55	9.04	83
Right superior parietal lobule	30	-55	56	8.93	130
Left cerebellum, posterior lobe, pyramis	-20	-62	-29	8.57	104
Left superior parietal lobule	-30	-57	54	8.29	28
Right cerebellum, posterior lobe, pyramis	30	-69	-31	8.19	36
Left cerebellum, posterior lobe, pyramis	-8	-73	-28	8.02	26
Right precuneus	18	-62	-27	7.96	74
Right SM1	22	-25	66	7.88	26

List of the brain regions showing a significant connectivity with the SMA (P < 0.05, corrected). The coordinates are given as stereotaxic coordinates referring to the atlas of Talairach and Tournoux (1988).

anti-phase movements, than normal controls (Johnson et al., 1998; Almeida et al., 2002; Ponsen et al., 2006). We used a slow movement rate (0.5 Hz) because it was easier for our patients to perform. It has been observed that both in- and anti-phase tasks are easily maintained in a stable rhythm at low frequencies, but anti-phase tasks often spontaneously convert to in-phase at higher frequencies (Kelso, 1984; Johnson et al., 1998). An external timing cue could help patients with Parkinson's disease to perform in-phase movement with more accuracy and stability, and with better coordination. However, for anti-phase movements, the external cue accentuated the tendency for patients to revert to in-phase movements (Johnson et al., 1998). The external cue may increase the complexity of bimanual tasks; patients have to perform the complex movement correctly and in time with the external cue. From our experience, patients with Parkinson's disease can perform movements at the required slow rate without external cues (Wu and Hallett, 2005). Therefore, we did not use an external timing cue in the current study. Since the rate of movement has a significant effect on brain activity (van Meter et al., 1995; Sadato et al., 1997a; Deiber et al., 1999), we gave all subjects sufficient time to practice the rate until they could perform it correctly. Actually, all patients could execute tasks properly at the required rate. There was no difference in the frequency of movements between groups. Since each subject's performance during fMRI was monitored by an investigator and recorded by video, we can assure that all subjects performed all tasks correctly and there was no phase error. Movement amplitude was controlled by asking the subjects to move the maximum amount possible for extension; whereas the amplitude of flexion was limited by the response device fixed to their hand. Therefore, it is unlikely that behavioural performance had obvious effects on the observed different brain activity or effective connectivity between groups.



Figure 6 (**A**) Brain areas more connected with the SMA-proper in patients with Parkinson's disease than in controls (upper row) and more connected in normal subjects than in patients with Parkinson's disease (lower row), during performance of in-phase movements; (**B**) Brain areas more connected with the SMA-proper in patients with Parkinson's disease than in controls (upper row) and more connected in normal subjects than in patients with Parkinson's disease (lower row), during performance of anti-phase movements. Results were thresholded at P < 0.05 (FWE corrected).

Changes in brain activity

Both patients and normal subjects had more activation in the SMA-proper, right SM1, cingulate motor area, premotor cortex and left cerebellum in performing bimanual movements compared with unimanual movement. An important finding is that the putamen and globus pallidus showed more activity in bimanual

movements compared with unimanual movement in controls, but not in patients with Parkinson's disease (Fig. 2A). This finding demonstrated that the basal ganglia could be more activated to perform bimanual tasks in healthy controls. In contrast, the dysfunction of basal ganglia appears not to allow further recruitment for the more complex bimanual movements in Parkinson's disease. We also found that the activation in the basal ganglia was decreased in patients with Parkinson's disease compared with controls during the performance of bimanual movements (Fig. 3). It has been suggested that the basal ganglia may be crucial in the neural control of bimanual coordination and may be specifically involved in the initiation phase of bimanual movements (Cardoso de Oliveira, 2002; Kraft *et al.*, 2007). Thus, it can certainly be possible that the damaged function of the basal ganglia may impair the ability of patients with Parkinson's disease to perform bimanual tasks.

Agreeing with previous findings (Johnson *et al.*, 1998; Serrien *et al.*, 2000; van den Berg *et al.*, 2000; Geuze, 2001; Almeida *et al.*, 2002; Ponsen *et al.*, 2006), our patients had more errors in performing anti-phase movements than in performing in-phase movements. To perform in-phase movements, homologous muscles are active simultaneously and symmetrically. For anti-phase movements, the homologous muscles are activated 180° out of phase; whereas contralateral antagonist muscles must move simultaneously. Performing anti-phase movement requires specific, sequential timing of muscle activation to maintain the required difference between the two hands, and is mirror-asymmetrical. In addition, attention needs to be maintained in order to keep the required phase relationship between the two hands. Thus, anti-phase movement is more complex than in-phase movement (Spencer and Ivry, 2007).

In both groups, several brain areas were more activated for anti-phase movements than for in-phase movements, including the premotor cortex, inferior frontal gyrus, postcentral gyrus and cerebellum. In contrast, the SMA-proper was more activated in anti-phase movements than in in-phase movements in controls (Sadato et al. 1997b; Toyokura et al., 1999, 2002; Immisch et al., 2001), but not in patients with Parkinson's disease. There was decreased activation in the SMA-proper in patients compared with controls in the performance of bimanual movements (Fig. 3). The hypoactivation of SMA secondary to dopamine deficiency in Parkinson's disease has been extensively reported in neuroimaging studies (Jenkins et al., 1992; Playford et al., 1992; Rascol et al., 1994; Jahanshani et al., 1995; Samuel et al., 1997; Haslinger et al., 2001; Buhmann et al., 2003). Our observation that the SMA-proper had more activation in bimanual movements than in unimanual movements suggests that the SMA could be more activated to perform bimanual movements in Parkinson's disease, at least at the early stage of the disorder. However, the activation in the SMA-proper could not be further increased in anti-phase movements than in in-phase movements, which indicates that there are no further resources in the SMA to be utilized during the more complex anti-phase movements in patients with Parkinson's disease. Some of our patients reverted anti-phase movement occasionally to in-phase movement. Studies on monkeys and humans with SMA damage have also revealed such a tendency to revert from mirror-asymmetrical to mirror-symmetrical movement (Luria, 1966; Brinkman, 1981; Chan and Ross, 1988). Given the crucial role of the SMA in bimanual coordination, we speculate that dysfunction of the SMA is likely to be an important contributor to the deficiency of patients with Parkinson's disease in performing bimanual movements, especially for anti-phase movements.

Besides the hypoactivation of the SMA and basal ganglia, we also observed hyperactivity in the SM1, premotor cortex, inferior frontal gyrus, superior parietal lobule, inferior parietal lobule, precuneus and cerebellum in patients with Parkinson's disease compared with controls in performing anti-phase movements (Fig. 3). All these regions have been suggested to have specific roles in bimanual coordination (Sadato et al., 1997b: Donchin et al., 1998, 2002; Toyokura et al., 1999; Kermadi et al., 2000; Tracy et al., 2001; de Jong et al., 2002; Iwamura et al., 2002; Meyer-Lindenberg et al., 2002; Ullen et al., 2003; Debaere et al., 2004; Wenderoth et al., 2004, 2005). The premotor cortex may have an important role in the higher control of bimanual coordination (Debaere et al., 2004), especially for anti-phase than for in-phase tasks (Sadato et al., 1997b; de Jong et al., 2002; Meyer-Lindenberg et al., 2002; Ullen et al., 2003). Wenderoth et al. (2005) found that the precuneus is more activated while performing bimanual movements than unimanual tasks, and attributed this to the more attention required for bimanual movements. The posterior lobe of the cerebellum is more specific to timing of more complex bilateral limb movements (Ullen et al., 2003). Blood flow increases in the anterior cerebellar vermis and hemisphere with increasing frequency for anti-phase coordination but not for the in-phase pattern (Meyer-Lindenberg et al., 2002). Furthermore, the cerebellum may specifically relate to the monitoring and correction of the spatiotemporal relationship between the limbs and its implementation into the required rhythm (Debaere et al., 2004).

The dysfunction of the basal ganglia and SMA should induce deterioration in performing bimanual tasks. However, after practice, our patients could execute the bimanual tasks at the same level as the normal subjects. Therefore, we speculate that the greater activity in the SM1, premotor cortex, parietal cortex, precuneus and cerebellum in our patients compared with controls is likely to provide the compensation for the dysfunction of the SMA and basal ganglia required in order to perform bimanual movements correctly (Rascol et al., 1997; Catalan et al., 1999; Sabatini et al., 2000; Wu and Hallett, 2005). In addition, we found that activations in the basal ganglia and SMA-proper were negatively correlated with UPDRS, whereas activations in the cerebellum, premotor cortex, parietal cortex and precuneus were positively correlated with UPDRS. These findings indicate that as the disorder progresses, dysfunction of the basal ganglia and SMA becomes more severe and contributions of these regions to the performance of bimanual movements may decrease. At the same time, the apparent compensatory effect in the cerebellum, premotor cortex, parietal cortex and precuneus is more significant.

The SM1, cerebellum, parietal cortex, precuneus and premotor cortex have been demonstrated to show increased activity as movements become more complex (Shibasaki *et al.*, 1993; Chen *et al.*, 1997; Catalan *et al.*, 1998; Ziemann and Hallett, 2001; Wu *et al.*, 2004; Verstynen *et al.*, 2005). Additionally, the cerebellum and premotor cortex were identified as the principal regions responding to the manipulation of complexity in bimanual coord-ination (Tracy *et al.*, 2001; Dabaere *et al.*, 2004). Thus, the greater activity in these regions may also partially be due to more difficulty in patients with Parkinson's disease than controls in performing bimanual tasks.

Changes in effective connectivity

In both groups, the SMA-proper connected with extensive areas during the performance of either in-phase or anti-phase tasks, such as the bilateral SM1, premotor cortex, basal ganglia and cerebellum (Fig. 4). These results indicate that brain motor networks, especially the regions related to bimanual coordination, are tightly connected in order to perform bimanual movements.

A significant difference between anti-phase and in-phase movements is that the connectivity between the left (dominant) SM1 and SMA-proper was increased, while the connectivity between the right (non-dominant) SM1 and SMA-proper was decreased in the anti-phase compared with in-phase conditions. An EEG study found that interhemispheric transmission of information is selectively driven between the bilateral SM1 as a function of task requirements, and bimanual movements are mainly controlled by the dominant hemisphere (Serrien et al., 2003). Using the Structural Equation Modelling method, Walsh et al. (2008) showed that the dominant hemisphere appears to initiate activity responsible for bimanual movement. Serrien et al. (2002) observed that repetitive transcranial magnetic stimulation of the SMA could impair temporal accuracy of bimanual movement performance, especially for anti-phase tasks, and suggested that the SMA has an important integrative role in the organization of bimanual configurations as a function of task complexity, and operates bilaterally with interhemispheric interactions adjusting the activity of both SM1. It is likely that the interregional interactions between the SMA and SM1 need to be shifted to the dominant side in order to perform the more complex anti-phase movement, which induces the observed different pattern of connectivity of the bilateral SM1 and SMA between anti-phase and in-phase movements.

Both groups also showed more connectivity with SMA-proper in some other regions, like the premotor cortex and precuneus during performing anti-phase movement than in performing in-phase movement (Table 5 and Fig. 5). These results suggest that to perform the more complex anti-phase movements, bimanual coordination-related brain areas should be more tightly connected.

During performing anti-phase movement, patients with Parkinson's disease had less connectivity in the left putamen and left dorsolateral prefrontal cortex compared with controls (Table 7 and Fig. 5). Using diffusion tensor imaging method, it has been shown that the SMA-proper connects to the putamen bilaterally (Lehéricy *et al.*, 2004). In Parkinson's disease, the dopamine uptake is mostly reduced in the putamen (Brooks *et al.*, 1990). Thus, the decreased connectivity between the putamen and SMA in Parkinson's disease is probably a consequence of the dysfunction of the basal ganglia. This, in turn, may disrupt the function of SMA and contribute to the difficulty in performing bimanual movements in Parkinson's disease.

The dorsolateral prefrontal cortex is specialized for 'attentionalcognitive' functions (Jueptner *et al.*, 1997; Yamasaki *et al.*, 2002). This region has been reported to be more activated for bimanual anti-phase tasks than for an in-phase task (Haslinger *et al.*, 2004). The connectivity between the dorsolateral prefrontal cortex and SMA was increased in anti-phase movements compared with the in-phase tasks in normal subjects, which suggests that a higher level of attention is required to carry out the demanding bimanual anti-phase movements (Haslinger et al., 2004). In contrast, the connectivity between the dorsolateral prefrontal cortex and SMA was not increased in the anti-phase condition in Parkinson's disease and was decreased compared with controls. A previous study reported that when performing movements that require attention to action, the connectivity between the prefrontal cortex and the premotor cortex was increased in healthy subjects but not in patients with Parkinson's disease (Rowe et al., 2002). Our previous study suggested that limited attentional resources might be a reason related to the difficulty in performing two tasks simultaneously in Parkinson's disease (Wu and Hallett, 2008). Thus, the abnormal connectivity between the dorsolateral prefrontal cortex and SMA may also be an indication of the disrupted attentional networks in Parkinson's disease, which is possibly a reason contributing to deteriorated bimanual coordination in performing anti-phase movements.

Patients with Parkinson's disease showed increased connectivity in the bilateral SM1, bilateral superior parietal lobule, right precuneus and bilateral cerebellum with SMA-proper in performance of anti-phase movements compared with controls (Table 7 and Fig. 5B). As these regions are involved in bimanual coordination, these increased connections are also likely to compensate for the defective basal ganglia to perform bimanual tasks correctly (Helmich *et al.*, 2009; Palmer *et al.*, 2009). However, the increased interregional interactions may also reflect a facet of the primary pathophysiology of Parkinson's disease, such as an inability to inhibit contextually inappropriate circuits secondary to abnormal basal ganglia outflow (Mink, 1996; Turner *et al.*, 2003; Grafton *et al.*, 2006).

The cerebellum and basal ganglia have distinct loops connecting with largely overlapping cortical areas (Middleton and Strick, 2000), and some motor control functions might be shared by basal ganglia and cerebellar motor systems (Desmurget et al., 2004). Both the basal ganglia and cerebellum regulate cortical excitability, but may have somewhat opposing influences (Liepert et al., 2004; Tamburin et al., 2004; Hallett, 2006). Thus, changed connectivity may also reflect a compensatory reaction of the cerebellum secondary to the damage of the basal ganglia. However, because the SMA receives significantly more basal ganglia input than cerebellar input (Akkal et al., 2007), even though the cerebellar compensation exists, it may not be strong enough to normalize the dysfunction in the SMA due to the damage of basal ganglia. We observed a significant increase of connectivity between the cerebellum and SMA in anti-phase than in in-phase movements in normal subjects but not in patients with Parkinson's disease. Possibly, the interaction between the cerebellum and SMA already achieves the limit in performing bimanual in-phase movements and could not be further increased to give more compensation for the more complex bimanual anti-phase movements. An alternative view is that the changes of the connectivity between the SMA and cerebellum may not always produce an adaptive response to restore normal motor function; it may be a part of the problem of motor deficits, rather than a solution (Grafton, 2004).

Brain 2010: 133; 2394–2409 | **2407**

In conclusion, our findings suggest that several reasons may contribute to the difficulty of performing bimanual anti-phase movements in patients with Parkinson's disease. The SMA and basal ganglia are hypoactivated. The pattern of interactions of neural networks is abnormal; the connectivity between the SMA and putamen is decreased. In addition, the attentional network is disrupted. The patients need to recruit more brain activity and increase connectivity in some brain regions, like the primary motor cortex and cerebellum to compensate for dysfunction of the basal ganglia and SMA in order to perform bimanual movements correctly.

Funding

The National Science Foundation of China (30870693); Ministry of Science and Technology (2006AA02A408); Intramural Research Program of the National Institute of Neurological Disorders and Stroke, National Institutes of Health (to M.H.).

References

- Akkal D, Dum R, Strick PL. Supplementary motor area and presupplementary motor area: targets of basal ganglia and cerebellar output. J Neurosci 2007; 27: 10659–73.
- Almeida QJ, Wishart LR, Lee TD. Bimanual coordination deficits with Parkinson's disease: the influence of movement speed and external cueing. Mov Disord 2002; 17: 30–7.
- Brinkman C. Lesions in supplementary motor area interfere with a monkey's performance of a bimanual coordination task. Neurosci Lett 1981; 27: 267–70.
- Brooks DJ, Ibanez V, Sawle GV, Quinn N, Lees AJ, Mathias CJ, et al. Differing patterns of striatal 18F-dopa uptake in Parkinson's disease, multiple system atrophy, and progressive supranuclear palsy. Ann Neurol 1990; 28: 547–55.
- Buhmann C, Glauche V, Sturenburg HJ, Oechsner M, Weiller C, Buchel C. Pharmacologically modulated fMRI - cortical responsiveness to levodopa in drug-naive hemiparkinsonian patients. Brain 2003; 126: 451–61.
- Cardoso de Oliveira S. The neuronal basis of bimanual coordination: recent neurophysiological evidence and functional models. Acta Psychol 2002; 110: 139–59.
- Catalan MJ, Honda M, Weeks RA, Cohen LG, Hallett M. The functional neuroanatomy of simple and complex sequential finger movements: a PET study. Brain 1998; 121: 253–64.
- Catalan MJ, Ishii K, Honda M, Samii A, Hallett M. A PET study of sequential finger movements of varying length in patients with Parkinson's disease. Brain 1999; 122: 483–95.
- Chan JL, Ross ED. Left-handed mirror writing following right anterior cerebral artery infarction: evidence for nonmirror transformation of motor programs by right supplementary motor area. Neurology 1988; 38: 59–63.
- Chen R, Gerloff C, Hallett M, Cohen LG. Involvement of the ipsilateral motor cortex in finger movements of different complexities. Ann Neurol 1997; 41: 247–54.
- Chen JT, Lin YY, Shan DE, Wu ZA, Hallett M, Liao KK. Effect of transcranial magnetic stimulation on bimanual movements. J Neurophysiol 2005; 93: 53–63.
- de Jong BM, Leenders KL, Paans AM. Right parieto-premotor activation related to limb-independent antiphase movement. Cereb Cortex 2002; 12: 1213–7.
- Debaere F, Wenderoth N, Sunaert S, Van Hecke P, Swinnen SP. Cerebellar and premotor function in bimanual coordination: parametric

neural responses to spatiotemporal complexity and cycling frequency. NeuroImage 2004; 21: 1416–27.

- Debaere F, Swinnen SP, Beatse E, Sunaert S, Van Hecke P, Duysens J. Brain areas involved in interlimb coordination: a distributed network. Neuroimage 2001; 14: 947–58.
- Deiber MP, Honda M, Ibanez V, Sadato N, Hallett M. Mesial motor areas in self-initiated versus externally triggered movements examined with fMRI: effect of movement type and rate. J Neurophysiol 1999; 81: 3065–77.
- Desmurget M, Grafton ST, Vindras P, Grea H, Turner RS. The basal ganglia network mediates the planning of movement amplitude. Eur J Neurosci 2004; 19: 2871–80.
- Donchin O, Gribova A, Steinberg O, Bergman H, Vaadia E. Primary motor cortex is involved in bimanual coordination. Nature 1998; 395: 274–8.
- Donchin O, Gribova A, Steinberg O, Mitz AR, Bergman H, Vaadia E. Single-unit activity related to bimanual arm movements in the primary and supplementary motor cortices. J Neurophysiol 2002; 88: 3498–517.
- Friston KJ, Frith CD, Frackowiak RS. Time-dependent changes in effective connectivity measured with PET. Hum Brain Mapp 1993b; 1: 69–80.
- Friston KJ, Frith CD, Liddle PF, Frackowiak RS. Functional connectivity: the principal component analysis of large (PET) data sets. J Cereb Blood Flow Metab 1993a; 13: 5–14.
- Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ. Psychophysiological and modulatory interactions in neuroimaging. NeuroImage 1997; 6: 218–29.
- Garraux G, McKinney C, Wu T, Kansanku K, Nolte G, Hallett M. Shared brain areas but not functional connections controlling movement timing and order. J Neurosci 2005; 25: 5290–7.
- Geuze RH. Stability of bimanual coordination in Parkinson's disease and cognitive modulation of intention. Motor Control 2001; 5: 361–84.
- Gitelman DR, Penny WD, Ashburner J, Friston K. Modeling regional and psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution. NeuroImage 2003; 19: 200–7.
- Grafton ST. Contributions of functional imaging to understanding parkinsonian symptoms. Curr Opin Neurobiol 2004; 14: 715–9.
- Grafton ST, Turner RS, Desmurget M, Bakay R, Delong M, Vitek J, et al. Normalizing motor-related brain activity: subthalamic nucleus stimulation in Parkinson disease. Neurology 2006; 66: 1192–99.
- Hallett M. Pathophysiology of dystonia. J Neural Transm Suppl 2006; 70: 485–88.
- Haslinger B, Erhard P, Altenmuller E, Hennenlotter A, Schwaiger M, Grafin von Einsiedel H, et al. Reduced recruitment of motor association areas during bimanual coordination in concert pianists. Hum Brain Mapp 2004; 22: 206–15.
- Haslinger B, Erhard P, Kampfe N, Boecker H, Rummeny E, Schwaiger M, et al. Event-related functional magnetic resonance imaging in Parkinson's disease before and after levodopa. Brain 2001; 124: 558–70.
- Helmich RC, Aarts E, de Lange FP, Bloem BR, Toni I. Increased dependence of action selection on recent motor history in Parkinson's disease. J Neurosci 2009; 29: 6105–13.
- Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. Neurology 1967; 17: 427-42.
- Immisch I, Waldvogel D, van Gelderen P, Hallett M. The role of the medial wall and its anatomical variations for bimanual antiphase and in-phase movements. NeuroImage 2001; 14: 674–84.
- Iwamura Y, Tanaka M, Iriki A, Taoka M, Toda T. Processing of tactile and kinesthetic signals from bilateral sides of the body in the postcentral gyrus of awake monkeys. Behav Brain Res 2002; 135: 185–90.
- Jahanshani M, Jenkins H, Brown RG, Marsden CD, Passingham RE, Brooks DJ. Self-initiated versus externally triggered movements. I. An investigation using measurement of regional cerebral blood flow with PET and movement-related potentials in normal and Parkinson's disease subjects. Brain 1995; 118: 913–33.

- Jenkins IH, Fernandez W, Playford ED, Lees AJ, Frackowiak RS, Passingham RE, et al. Impaired activation of the supplementary motor area in Parkinson's disease is reversed when akinesia is treated with apomorphine. Ann Neurol 1992; 32: 749–57.
- Johnson KA, Cunnington R, Bradshaw JL, Phillips JG, Iansek R, Rogers MA. Bimanual co-ordination in Parkinson's disease. Brain 1998; 121: 743–53.
- Jueptner M, Stephan KM, Frith CD, Brooks DJ, Frackowiak RSJ. Anatomy of motor learning. I. Frontal cortex and attention to action. J Neurophysiol 1997; 77: 1313–24.
- Kelso JA. Phase transitions and critical behavior in human bimanual coordination. Am J Physiol 1984; 246: R1000-4.
- Kermadi I, Liu Y, Rouiller EM. Do bimanual motor actions involve the dorsal premotor (PMd), cingulate (CMA) and posterior parietal (PPC) cortices? Comparison with primary and supplementary motor cortical areas. Somatosens Motor Res 2000; 17: 255–71.
- Kraft E, Chen AW, Flaherty AW, Blood AJ, Kwong KK, Jenkins BG. The role of the basal ganglia in bimanual coordination. Brain Res 2007; 1151: 62–73.
- Lang AE, Fahn S. Assessment of Parkinson's disease. In: Munsat TL, editor. Quantification of neurological deficit. Boston: Butterworths; 1989. p. 285–309.
- Lehéricy S, Ducros M, Krainik A, Francois C, Van de Moortele P, Ugurbil K, et al. 3-D diffusion tensor axonal tracking shows distinct SMA and Pre-SMA projections to the human striatum. Cereb Cortex 2004; 14: 1302–9.
- Liepert J, Kucinski T, Tuscher O, Pawlas F, Baumer T, Weiller C. Motor cortex excitability after cerebellar infarction. Stroke 2004; 35: 2484–88.
- Luria AR. Higher cortical functions in man. New York: Basic Books; 1966.
- Ma Z, Wang ZJ. Dynamic analysis of probabilistic boolean network for fMRI study in Parkinson's disease. Conf Proc IEEE Eng Med Biol Soc 2008; 2008: 161–4.
- Meyer-Lindenberg A, Ziemann U, Hajak G, Cohen L, Berman KF. Transitions between dynamical states of differing stability in the human brain. Proc Natl Acad Sci USA 2002; 99: 10948–53.
- Middleton FA, Strick PL. Basal ganglia and cerebellar loops: motor and cognitive circuits. Brain Res Brain Res Rev 2000; 31: 236–50.
- Mink J. The basal ganglia: focused selection and inhibition of competing motor programs. Prog Neurobiol 1996; 50: 381–425.
- Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. Neuropsychologia 1971; 9: 97–113.
- Palmer SJ, Eigenraam L, Hoque T, McCaig RG, Troiano A, McKeown MJ. Levodopa-sensitive, dynamic changes in effective connectivity during simultaneous movements in Parkinson's disease. Neuroscience 2009; 158: 693–704.
- Playford ED, Jenkins IH, Passingham RE, Nutt J, Frackowiak RS, Brooks DJ. Impaired mesial frontal and putamen activation in Parkinson's disease: a positron emission tomography study. Ann Neurol 1992; 32: 151–61.
- Ponsen MM, Daffertshofer A, van den Heuvel E, Wolters ECh, Beek PJ, Berendse HW. Bimanual coordination dysfunction in early, untreated Parkinson's disease. Parkinsonism Relat Disord 2006; 12: 246–52.
- Rascol O, Sabatini U, Chollet F, Fabre N, Senard JM, Montastruc JL, et al. Normal activation of the supplementary motor area in patients with Parkinson's disease undergoing long-term treatment with levodopa. J Neurol Neurosurg Psychiatry 1994; 57: 567–71.
- Rascol O, Sabatini U, Fabre N, Brefel C, Loubinoux I, Celsis P, et al. The ipsilateral cerebellar hemisphere is overactive during hand movements in akinetic parkinsonian patients. Brain 1997; 120: 103–10.
- Rowe J, Stephan KE, Friston K, Frackowiak R, Lees A, Passingham R. Attention to action in Parkinson's disease: impaired effective connectivity among frontal cortical regions. Brain 2002; 125: 276–89.
- Sabatini U, Boulanouar K, Fabre N, Martin F, Carel C, Colonnese C, et al. Cortical motor reorganization in akinetic patients with Parkinson's disease: a functional MRI study. Brain 2000; 123: 394–403.

- Sadato N, Yonekura Y, Waki A, Yamada H, Ishii Y. Role of the supplementary motor area and the right premotor cortex in the coordination of bimanual finger movements. J Neurosci 1997b; 17: 9667–74.
- Sadato N, Ibanez V, Campbell G, Deiber MP, Le Bihan D, Hallett M. Frequency-dependent changes of regional cerebral blood flow during finger movements: functional MRI compared to PET. J Cereb Blood Flow Metab 1997a; 17: 670–9.
- Samuel M, Ceballos-Baumann AO, Blin J, Uema T, Boecker H, Passingham RE, et al. Evidence for lateral premotor and parietal overactivity in Parkinson's disease during sequential and bimanual movements. A PET study. Brain 1997; 120: 963–76.
- Serrien D, Cassidy MJ, Brown P. The importance of the dominant hemisphere in the organization of bimanual movements. Hum Brain Mapp 2003; 18: 296–305.
- Serrien D, Strens LHA, Oliviero A, Brown P. Repetitive transcranial magnetic stimulation of the supplementary motor area (SMA) degrades bimanual movement control in humans. Neurosci Lett 2002; 328: 89–92.
- Serrien DJ, Steyvers M, Debaere F, Stelmach GE, Swinnen SP. Bimanual coordination and limb-specific parameterization in patients with Parkinson's disease. Neuropsychologia 2000; 38: 1714–22.
- Shibasaki H, Sadato N, Lyshkow H, Yonekura Y, Honda M, Nagamine T, et al. Both primary motor cortex and supplementary motor area play an important role in complex finger movement. Brain 1993; 116: 1387–98.
- Spencer RMC, Ivry RB. The temporal representation of in-phase and anti-phase movements. Hum Mov Sci 2007; 26: 226–34.
- Stephan KE, Marshall JC, Friston KJ, Rowe JB, Ritzl A, Zilles K, et al. Lateralized cognitive processes and lateralized task control in the human brain. Science 2003; 301: 384–6.
- Stephan KM, Binkofski F, Halsband U, Dohle C, Wunderlich G, Schnitzler A, et al. The role of ventral medial wall motor areas in bimanual co-ordination: a combined lesion and activation study. Brain 1999; 122: 351–68.
- Swinnen SP. Intermanual coordination: from behavioural principles to neural network interactions. Nat Rev Neurosci 2002; 3: 348–59.
- Talairach J, Tournoux P. Co-Planar Stereotaxic Atlas of the Human Brain. Thieme Verlag, Stuttgart. 1988.
- Tamburin S, Fiaschi A, Marani S, Andreoli A, Manganotti P, Zanette G. Enhanced intracortical inhibition in cerebellar patients. J Neurol Sci 2004; 217: 205–10.
- Tanji J, Hoshi E. Behavioral planning in the prefrontal cortex. Curr Opin Neurobiol 2001; 11: 164–70.
- Toyokura M, Muro I, Komiya T, Obara M. Relation of bimanual coordination to activation in the sensorimotor cortex and supplementary motor area: analysis using functional magnetic resonance imaging. Brain Res Bull 1999; 48: 211–7.
- Toyokura M, Muro I, Komiya T, Obara M. Activation of presupplementary motor area (SMA) and SMA proper during unimanual and bimanual complex sequences: an analysis using functional magnetic resonance imaging. J Neuroimaging 2002; 12: 172–8.
- Tracy JI, Faro SS, Mohammed FB, Pinus AB, Madi SM, Laskas JW. Cerebellar mediation of the complexity of bimanual compared to unimanual movements. Neurology 2001; 57: 1862–9.
- Tuller B, Kelso JA. Environmentally-specified patterns of movement coordination in normal and split-brain subjects. Exp Brain Res 1989; 75: 306–16.
- Turner RS, Grafton ST, McIntosh AR, DeLong MR, Hoffman JM. The functional anatomy of parkinsonian bradykinesia. Neuroimage 2003; 19: 163–79.
- Ullen F, Forssberg H, Ehrsson HH. Neural networks for the coordination of the hands in time. J Neurophysiol 2003; 89: 1126–35.
- van den Berg C, Beek PJ, Wagenaar RC, van Wieringen PCW. Coordination disorders in patients with Parkinson's disease: a study of paced rhythmic forearm movements. Exp Brain Res 2000; 134: 174–86.

- van Eimeren T, Monchi O, Ballanger B, Strafella AP. Dysfunction of the default mode network in Parkinson disease: a functional magnetic resonance imaging study. Arch Neurol 2009; 66: 877–83.
- van Meter JW, Maisog JM, Zeffiro TA, Hallett M, Herscovitch P, Rapoport SI. Parametric analysis of functional neuroimages: application to a variable-rate motor task. NeuroImage 1995; 2: 273–83.
- Verstynen T, Diedrichsen J, Albert N, Aparicio P, Ivry RB. Ipsilateral motor cortex activity during unimanual hand movements relates to task complexity. J Neurophysiol 2005; 93: 1209–22.
- Walsh RR, Small SL, Chen EE, Solodkin A. Network activation during bimanual movements in humans. NeuroImage 2008; 43: 540–53.
- Wenderoth N, Debaere F, Sunaert S, Van Hecke P, Swinnen SP. Parieto-premotor areas mediate directional interference during bimanual movements. Cereb Cortex 2004; 14: 1153–63.
- Wenderoth N, Debaere F, Sunaert S, Swinnen SP. The role of anterior cingulate cortex and precuneus in the coordination of motor behaviour. Eur J Neurosci 2005; 22: 235–46.

- Wu T, Hallett M. A functional MRI study of automatic movements in patients with Parkinson's disease. Brain 2005; 128: 2250–9.
- Wu T, Hallett M. Neural correlates of dual task performance in patients with Parkinson's disease. J Neurol Neurosurg Psychiatry 2008; 79: 760–6.
- Wu T, Kansaku K, Hallett M. How self-initiated memorized movements become automatic: a fMRI study. J Neurophysiol 2004; 91: 1690–8.
- Wu T, Chan P, Hallett M. Effective connectivity of neural networks in automatic movements in Parkinson's disease. NeuroImage 2010; 49: 2581–7.
- Yamasaki H, LaBar KS, McCarthy G. Dissociable prefrontal brain systems for attention and emotion. Proc Natl Acad Sci USA 2002; 99: 11447–51.
- Ziemann U, Hallett M. Hemispheric asymmetry of ipsilateral motor cortex activation during unimanual motor tasks: further evidence for motor dominance. Clin Neurophysiol 2001; 112: 107–13.